

The Geometry of Grauert Tubes and Complexification of Symmetric Spaces

D. Burns* S. Halverscheid R. Hind

February 1, 2008

Abstract

We study the canonical complexifications of non-compact Riemannian symmetric spaces by the Grauert tube construction. We determine the maximal such complexification, a domain already constructed by Akhiezer and Gindikin [1], and show that this domain is Stein. We also determine when invariant complexifications, including the maximal one, are Hermitian symmetric. This is expressed simply in terms of the ranks of the symmetric spaces involved.

1 Introduction

The purpose of this paper is to investigate certain canonical complexifications of irreducible Riemannian symmetric spaces of the noncompact type. We will be interested in determining whether these complexifications are rigid in the sense that their complex automorphism group is isomorphic to the isometry group of the symmetric space (this is always a subgroup) or whether the complex manifold exhibits additional symmetries. We will also deal with a question of

*Partially supported by NSF, DMS-0104047.

Steinness, verifying a conjecture of D. Akhiezer and S. Gindikin.

The construction of the canonical complexifications is actually very general and applies to any real-analytic Riemannian manifold. A general rigidity result was proven in our previous paper [6] for the complex manifolds associated to compact Riemannian manifolds, but almost nothing is known about the complexifications of arbitrary noncompact manifolds. In the case of symmetric spaces we are continuing work of S-J. Kan and D. Ma from [15].

We start with the following theorem proved independently by V. Guillemin and M. Stenzel in [7] and by L. Lempert and R. Szöke in [17].

Suppose that (M, g) is a real-analytic Riemannian manifold of dimension n . Identify M with the zero section in TM . Let $\rho : TM \rightarrow \mathbb{R}$ be the length, with respect to g , of tangent vectors.

Theorem 1 *There exists a unique complex structure on all sufficiently small neighbourhoods of M in TM such that the following conditions hold.*

- (i) ρ^2 is strictly plurisubharmonic and the corresponding Kähler metric restricts to g on M .
- (ii) ρ is a solution of the homogeneous complex Monge-Ampère equation $(dd^c \rho)^n = 0$ on $T^r M \setminus M$, where $M \subset T^r M$.

Let $T^r M = \{v \in TM | \rho(v) < r\}$. Then when M is a compact manifold or a symmetric space the complex structure exists on the disk bundles $T^r M$ for r sufficiently small. In this case we call the resulting complex manifolds $T^r M$ Grauert tubes, and r will often be called the radius of the tube. We will also call the complex manifolds Grauert domains even if the underlying subset of $T^r M$ is not necessarily a tube. For a general noncompact M the canonical complex structure need not exist on any $T^r M$.

An equivalent characterization of these complex manifolds (see also [17]) is that the complex structure is the unique one making the leaves of the Riemann foliation with their natural complex structures into holomorphic curves. In other

words, for any geodesic $\gamma : \mathbb{R} \rightarrow M$ the map

$$d\gamma : \mathbb{C} \rightarrow TM$$

$$s + it \mapsto (\gamma(s), t\gamma'(s))$$

is actually holomorphic with respect to the Grauert domain structure, wherever that structure is defined.

We remark here that all Grauert tubes have an antiholomorphic involution $\sigma : T^r M \rightarrow T^r M$ given by $v \mapsto -v$.

By functoriality of the construction, the differential of any isometry of M is actually a biholomorphism of a Grauert tube $T^r M$. We will say that a Grauert domain N , invariant under the action of $Isom(M, g)$, is *rigid* if its complex automorphism group $Aut_{\mathbb{C}}(X) = dIsom(M, g)$.

The following theorem is taken from [6].

Theorem 2 *Any Grauert tube $T^r M$ of finite radius r over a compact manifold M is rigid.*

Now, for symmetric (or locally symmetric) spaces of nonnegative curvature the canonical complex structure exists on the whole tangent bundle TM (see for instance [22]). The resulting complex manifolds for compact rank one spaces are described by G. Patrizio and P-M. Wong in [19]. As a particular case, the Grauert tube complex structure on TS^2 associated to the standard round metric on S^2 is biholomorphic to the complex quadric $\{z_1^2 + z_2^2 + z_3^2 = 1\}$ in $\mathbb{C}^3$. Thus such tubes of infinite radius are not rigid in general.

For Riemannian manifolds with some negative sectional curvature, the whole tangent bundle cannot be given the structure of a Grauert domain. In fact the following theorem of Lempert and Szöke (see [17] again) gives an upper bound on the radius of a Grauert tube.

Theorem 3 *Suppose that the sectional curvature of a 2-plane of (M, g) is equal to $-\lambda$, with $\lambda > 0$. Then if a Grauert tube structure exists on $T^r M$ the radius*

satisfies

$$r \leq \frac{\pi}{2\sqrt{-\lambda}}.$$

Hence for manifolds (M, g) with some negative sectional curvature there is a finite radius r_{max} for which a Grauert tube structure exists on $T^r M$ if and only if $r \leq r_{max}$. We emphasize that Theorem 2 holds even if $r = r_{max}$.

From now on we will concentrate on the case when (M, g) is a symmetric space of the noncompact type. The following theorem is proven by Kan and Ma in [15].

Theorem 4 *Let (M, g) be a symmetric space and $T^r M$ an associated Grauert tube with $r < r_{max}$. Then $T^r M$ is either rigid or the ball.*

It turns out that there is at most one r for which $T^r M$ is covered by a ball, so we can say that all but at most one $T^r M$ are rigid for $r < r_{max}$. One surprising result of this paper is that for rank one symmetric spaces of noncompact type, the tubes $T^{r_{max}} M$ are never rigid, in fact they are all Hermitian symmetric.

We define the *maximal Grauert domain* associated to a real-analytic (M, g) to be the largest connected domain in TM containing M on which our canonical complex structure exists, that is, on which we can define a complex structure satisfying the conditions of Theorem 1.

Without loss of generality suppose that (M, g) is an irreducible symmetric space of noncompact type. It is not hard to show that when (M, g) is of rank one, the maximal Grauert domain is just the Grauert tube of maximal radius. However for higher rank cases the maximal Grauert domain is always larger.

In section 2 we will show that the maximal Grauert domains can be described algebraically, and correspond to the domains defined and studied by D. Akhiezer and S. Gindikin amongst others, see [1].

Let M be written in Klein form as $M = G/K$ where G is a connected semisimple Lie group with finite center and K is a maximal compact subgroup.

The groups G and K can be complexified to linear algebraic groups $G^{\mathbb{C}}$ and $K^{\mathbb{C}}$ over $\mathbb{C}$. Let $o = e.K^{\mathbb{C}} \in G^{\mathbb{C}}/K^{\mathbb{C}}$.

Now let $\mathfrak{g}$ be the Lie algebra of G and $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ be a Cartan decomposition corresponding to the pair (G, K) . We let $\mathfrak{a}$ be a maximal abelian subalgebra of $\mathfrak{p}$ and Σ be a root system of $\mathfrak{g}$ with respect to $\mathfrak{a}$.

We will identify $G(o)$ with $M = G/K \hookrightarrow G^{\mathbb{C}}/K^{\mathbb{C}}$. Each G -orbit in $G^{\mathbb{C}}/K^{\mathbb{C}}$ intersects $\exp(i\mathfrak{a}).o$ in an orbit of the Weyl group.

Finally, following [1] we define

$$\omega = \{H \in \mathfrak{a} \mid |\alpha(H)| < \frac{\pi}{2} \forall \alpha \in \Sigma\}.$$

Theorem 5 *The maximal Grauert domain associated to $M = G/K$ is biholomorphic to the domain $D = G(\exp(i\omega)).o \subset G^{\mathbb{C}}/K^{\mathbb{C}}$.*

This is essentially contained in [1] or [22] but we clarify it in section 2.

There is an alternative for the complex automorphism group of maximal Grauert domains. We prove that the domains are either rigid or biholomorphic to Hermitian symmetric spaces. In fact we will prove the following.

Theorem 6 *Let $\omega' \subseteq \omega$ be a symmetric convex domain invariant under the action of the Weyl group. Then the domain $D' = G(\exp(i\omega')).o$ is either Hermitian symmetric or $\text{Aut}_{\mathbb{C}}(D') \cong \text{Isom}(M)$.*

It remains to decide in specific cases whether a maximal Grauert domain is rigid or Hermitian symmetric. In the second case, we will have a Hermitian symmetric space with an antiholomorphic involution whose fixed point set is isomorphic to the original symmetric space. Such objects have been classified by H. Jaffee, see [10] and [11]. An immediate consequence is that the maximal Grauert domain of $SL(3, \mathbb{R})/SO(3)$ is rigid.

Suppose then that we have a Hermitian symmetric space N with an antiholomorphic involution σ whose fixed point set is our symmetric space $M =$

G/K . Then N is always a Grauert domain corresponding to M . We need to decide whether N is the maximal domain. Suppose $N = G'/K'$ in Klein form and $\mathfrak{g}' = \mathfrak{t}' + \mathfrak{p}'$ is its Cartan decomposition. Again letting $\mathfrak{g} = \mathfrak{t} + \mathfrak{p}$ be the decomposition of the Lie algebra of G , since $M \subset N$ is totally geodesic we have $\mathfrak{t} \subset \mathfrak{t}'$ and $\mathfrak{p} \subset \mathfrak{p}'$. Let $\mathfrak{a}$ be a maximal abelian subalgebra of $\mathfrak{p}$ and $\mathfrak{a}'$ with $\mathfrak{a} \subset \mathfrak{a}' \subset \mathfrak{p}'$ be a maximal abelian subalgebra of $\mathfrak{p}'$. Let $\Sigma \subset \mathfrak{a}^*$ be the root system of $\mathfrak{g}$ with respect to $\mathfrak{a}$ and $\Sigma' \subset \mathfrak{a}'^*$ be the root system of $\mathfrak{g}'$ with respect to $\mathfrak{a}'$.

Theorem 7 *The Hermitian space N is the maximal Grauert domain corresponding to M if and only if*

$$\max_{\alpha \in \Sigma} |\alpha(H)| = \max_{\alpha \in \Sigma'} |\alpha(H)| \forall H \in \mathfrak{a}.$$

For example, if M itself is Hermitian symmetric, then the maximal Grauert domain is biholomorphic to the product $M \times \overline{M}$ into which M embeds diagonally. A result close to this special case of the theorem is contained in [27].

The condition in Theorem 8 is sometimes awkward to check, especially in exceptional cases, but fortunately it is equivalent to the simple condition given in the next proposition.

Proposition 8 *The Hermitian space N is the maximal Grauert domain corresponding to M if and only if $\text{rank}(N) = 2\text{rank}(M)$.*

Combining Proposition 8 with H. Jaffee's classification, it is easy to obtain a general description of the maximal Grauert domains of irreducible symmetric spaces of the noncompact type. This section will conclude with the resulting list.

Whether they are Hermitian symmetric or not, we are able to prove the following general result on the structure of the maximal Grauert domains.

Theorem 9 *The maximal Grauert domain associated to a symmetric space is Stein.*

This result was conjectured by D. Akhiezer and S. Gindikin in [1], who showed some examples of this property. These domains are intimately related to the linear cycle spaces of P.A. Griffiths, see [26] for a review. Several examples of the theorem were thus shown by Huckleberry, Wolf and Zireau in various papers on linear cycle spaces (see [26] for references), using the fact that the linear cycle spaces are Stein, see [25], for example. In particular, the results of the current paper were in part motivated by the appearance of “hidden symmetries” for some linear cycle spaces, as pointed out to us by Joe Wolf.

The proof of the last theorem is by a direct characterization of the G -invariant plurisubharmonic functions on the maximal Grauert domain. Earlier examples of this result are due to Azad and Loeb [3] for compact symmetric spaces, and K.-H. Neeb [18] for certain non-degenerate semigroups.

After giving the algebraic description of maximal Grauert domains in section 2, we show in section 3 that for a Hermitian symmetric space the corresponding maximal domain is simply the product of the space with itself. This is used to prove Theorem 7 in section 4. In section 5 the maximality condition is shown to be equivalent to the simpler statement in Proposition 8. In section 6 we prove the alternative for the biholomorphism groups of the Grauert domains. Finally in section 7 we prove that the maximal Grauert domains are Stein by characterizing, as already noted, their G -invariant plurisubharmonic functions.

The third author would like to thank Brian Hall for several enlightening discussions on these topics.

Maximal Grauert domains of noncompact symmetric spaces

$SL(n, \mathbb{R})/SO(n), n > 2$	rigid
$SU^*(2n)/Sp(n)$	rigid
$SU(p, q)/S(U_p \times U_q)$	product
$SO_0(2, 1)/SO(2)$	product
$SO_0(p, 1)/SO(p), p > 2$	$SO_0(p, 2)/SO(p) \times SO(2)$
$SO_0(p, 2)/SO(p) \times SO(2)$	product
$SO_0(p, q)/SO(p) \times SO(q), q > 2$	rigid
$SO^*(2n)/U(n)$	product
$Sp(n, \mathbb{R})/U(n)$	product
$Sp(p, q)/Sp(p) \times Sp(q)$	$SU(2p, 2q)/S(U_{2p} \times U_{2q})$
$SL(n, \mathbb{C})/SU(n), n > 2$	rigid
$SO(n, \mathbb{C})/SO(n), n > 3$	rigid
$Sp(n, \mathbb{C})/Sp(n), n > 1$	rigid
$(\mathfrak{e}_{6(-14)}, \mathfrak{so}(10) + \mathbb{R})$	product
$(\mathfrak{e}_{7(-25)}, \mathfrak{e}_6 + \mathbb{R})$	product
$(\mathfrak{f}_{4(-20)}, \mathfrak{so}(9))$	$(\mathfrak{e}_{6(-14)}, \mathfrak{so}(10) + \mathbb{R})$
all other exceptional spaces	rigid

The notation above is taken from the book [9]. We notice that all rank 1 examples have Hermitian symmetric maximal Grauert tubes. Some real symmetric spaces do appear as the fixed point sets of involutions on Hermitian symmetric spaces but nevertheless have rigid maximal domains. Examples of this are the spaces $SO_0(p, q)/SO(p) \times SO(q)$, for $q > 2$ which appear as the fixed point sets of involutions on $SU(p, q)/S(U_p \times U_q)$. However, if p and q are even, then $SU(p, q)/S(U_p \times U_q)$ is a maximal Grauert domain. Another similar example is $SU^*(8)/Sp(4)$ inside $(\mathfrak{e}_{7(-25)}, \mathfrak{e}_6 + \mathbb{R})$.

2 Algebraic description of maximal Grauert domains

For a noncompact symmetric space (M, g) , let $G, K, G^{\mathbb{C}}, K^{\mathbb{C}}, \mathfrak{g}, \mathfrak{k}, \mathfrak{p}, \mathfrak{a}$ be as in the introduction, that is, $M = G/K$ where G has Lie algebra $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ and $\mathfrak{a} \subset \mathfrak{p}$ is maximal abelian. Let Σ be a root system of $\mathfrak{g}$ with respect to $\mathfrak{a}$ for the decomposition

$$\mathfrak{g} = Z(\mathfrak{a}) \oplus \bigoplus_{\alpha \in \Sigma} \mathfrak{g}^{\alpha} \quad (1)$$

of the adjoint representation of $\mathfrak{a}$. Here, $Z(\mathfrak{a})$ denotes the center of $\mathfrak{a}$.

We can identify the tangent bundle TM with $G \times_K \mathfrak{p}$ where for $k \in K, g \in G$ and $X \in \mathfrak{p}$,

$$k(g, X) = (gk^{-1}, Ad(k)X).$$

With this identification $G \times_K (Ad(K)\omega)$ is a subdomain of TM containing M , where

$$\omega = \{H \in \mathfrak{a} \mid |\alpha(H)| < \frac{\pi}{2} \forall \alpha \in \Sigma\}.$$

As in the introduction, let $D = G.exp(i\omega).o \subset G^{\mathbb{C}}/K^{\mathbb{C}}$.

Then it is proven in [1] that the map

$$\phi : G \times_K (Ad(K)\omega) \rightarrow D$$

$$(g, Ad(k)H) \mapsto gk \exp(iH).o$$

is a real-analytic G -equivariant diffeomorphism.

Furthermore, the leaves of the Riemann foliation map onto holomorphic curves in D and so the pull-back of the complex structure on D gives $G \times_K (Ad(K)\omega)$ the (unique) adapted complex structure.

It remains to show that this is actually the maximal domain. To do this, suppose that $H \in \omega$ and that there exists some $\alpha \in \Sigma$ with $\alpha(H) = \frac{\pi}{2}$. Let γ be the geodesic in $M = G/K$ with $\gamma'(0) = H$. Then we have a map

$$d\gamma : \mathbb{C} \rightarrow TM$$

$$s + it \mapsto (\gamma(s), t\gamma'(s)).$$

We need to show that a Grauert domain structure can extend only over $d\gamma(|t| < 1)$. In [22], section 3, a precise criterion describes how far the complex structure can be extended for symmetric spaces. In the following, this criterion will be related to the root decomposition (1).

Note that the linear operator on $\mathfrak{p}$

$$Y \mapsto R(Y, H)H := -(\text{ad}H)^2(Y)$$

is the Jacobi operator for the geodesic γ in $T_{eK}G/K \cong \mathfrak{p}$. Since $Z(\mathfrak{a}) \cap \mathfrak{p} = \mathfrak{a}$, the decomposition into eigenspaces of the Jacobi operator is given by

$$\mathfrak{p} = \mathfrak{a} \oplus \sum_{\alpha \in \Sigma} \{w - \theta(w) | w \in \mathfrak{g}^\alpha\}.$$

In case of a Riemannian symmetric space of non-compact type, the eigenvalues $-\alpha(H)^2$ of $-(\text{ad}H)^2$ are non-positive.

Since the Riemann curvature tensor is parallel, the solutions of the Jacobi equation

$$Y'' = -R(Y, \gamma')\gamma'$$

are given by $t \mapsto f_j(t)v_j(t)$, where the f_j are functions and the v_j are the parallelly transported vector fields along γ such that $v_j(0)$ is an eigenvector of the Jacobi operator. For an eigenvector $v_j = w_j - \theta(w_j) \in \mathfrak{p}$ of $-(\text{ad}H)^2$, where $w_j \in \mathfrak{g}_{\alpha_j}$, with eigenvalue $\alpha_j(H) < 0$, one obtains the fundamental solutions $\cosh(\alpha_j(H)t)$ and $\sinh(\alpha_j(H)t)$ for f_j . It is understood that a root α can appear repeatedly here. For v_j with eigenvalue 0, the fundamental solutions are obviously 1 and t . An eigenbasis $v_1, \dots, v_n$ in $\mathfrak{p}$ of $-(\text{ad}H)^2$ determines a basis

$$\{Y_j^{\text{hor}} \cosh(\alpha_j(H)t)v_j(t), Y_j^{\text{ver}} \sinh(\alpha_j(H)t)v_j(t), j = 1, \dots, n\}$$

for Jacobi fields along γ . These have the properties $Y_j^{\text{ver}}(0) = 0$ and $\nabla Y_j^{\text{hor}}(0) = 0$, $Y_j^{\text{hor}}(0) = \nabla Y_j^{\text{ver}}(0) = v_j$.

Jacobi fields along γ are in 1-1 correspondence to vector fields along $d\gamma$ invariant under the geodesic flow and the fibre multiplication $N_\sigma : TM \rightarrow TM, v \mapsto \sigma \cdot v$ for $\sigma \in \mathbb{R}$. The fields ξ_j corresponding to Y_j^{hor} and η_j corresponding to Y_j^{ver} are a frame of $T(TM)$ along $d\gamma|_{\mathbb{C} \setminus \mathbb{R}}$. In [22], it is shown that the almost complex tensor for the adapted complex structure along $d\gamma$ w. r. t. the frame $\xi_1, \eta_1, \xi_2, \eta_2, \dots, \xi_n, \eta_n$ is given by n blocks. A block for v_j with non-zero eigenvalue is given by

$$J_j = \begin{pmatrix} -\frac{\text{Reg}_j}{\text{Img}_j} & -\text{Img}_j - \frac{(\text{Reg}_j)^2}{\text{Img}_j} \\ \frac{1}{\text{Img}_j} & \frac{\text{Reg}_j}{\text{Img}_j} \end{pmatrix}$$

where $g_j(t + is) = \frac{1}{\alpha_j(H)} \tanh((t + is)\alpha_j(H))$. Inspection of the J_j shows that there are poles at sH if and only if $\alpha_j(sH) \in \frac{\pi}{2}\mathbb{Z}$ for some α_j . The blocks for zero-eigenvalues do not contribute poles. Note that the poles at $s = 0$ are due to the fact that the $\xi_1, \eta_1, \xi_2, \eta_2, \dots, \xi_n, \eta_n$ are not a frame there. This confirms the result.

3 Maximal Grauert domain of a Hermitian symmetric space of noncompact type

Let $M = G/K$ be Hermitian symmetric of non-compact type. Thus M can be thought of as a bounded domain in some $\mathbb{C}^n$. More precisely, there is an open embedding of $M = G/K$ in its compact dual (Hermitian) symmetric space U/K . Let $\overline{M}$ denote M with the opposite complex structure. Then M embeds diagonally in $M \times \overline{M}$ and is the fixed point set of the antiholomorphic involution $(x, y) \mapsto (y, x)$. It is the aim of this section to show that $M \times \overline{M}$ is actually the maximal Grauert domain corresponding to M . We will follow very closely Chapter VIII from the book [9].

We use similar notation to the introduction, but now let $\mathfrak{g}^{\mathbb{C}}$, $\mathfrak{t}^{\mathbb{C}}$ and $\mathfrak{p}^{\mathbb{C}}$ denote the complexifications of $\mathfrak{g}$, $\mathfrak{t}$ and $\mathfrak{p}$ respectively. Also, let $\mathfrak{h}$ be a maximal

abelian subalgebra of $\mathfrak{t}$, then the complexification $\mathfrak{h}^{\mathbb{C}}$ is a Cartan subalgebra of $\mathfrak{g}^{\mathbb{C}}$. Let Δ be the non-zero roots of $\mathfrak{g}^{\mathbb{C}}$ with respect to $\mathfrak{h}^{\mathbb{C}}$ and $\{H_\alpha | \alpha \in \Delta\}$ be the elements of $\mathfrak{h}^{\mathbb{C}}$ uniquely defined so that $\alpha(H) = B(H, H_\alpha)$ for all $H \in \mathfrak{h}^{\mathbb{C}}$, where B is the Killing form. We let $\mathfrak{g}^{\mathbb{C}\alpha}$ be the corresponding root subspaces.

Now, there exists a collection $\Gamma \subset \Delta$ of strongly orthogonal roots $\Gamma = \{\gamma_1, \dots, \gamma_r\}$ and corresponding vectors $X_\gamma \in \mathfrak{g}^{\mathbb{C}\gamma}$ such that

$$\mathfrak{a} = \sum_{\gamma \in \Gamma} \mathbb{R}(X_\gamma + X_{-\gamma})$$

is a maximal abelian subspace of $\mathfrak{p}$. The restricted roots Σ in the decomposition (1) are given by (case C_r):

$$\Sigma = \{\pm \frac{\gamma_i + \gamma_j}{2}, 1 \leq j \leq i \leq r\} \cup \{\pm \frac{\gamma_i - \gamma_j}{2}, 1 \leq j < i \leq r\},$$

or (case BC_r):

$$\Sigma = \{\pm \frac{\gamma_i + \gamma_j}{2}, 1 \leq j \leq i \leq r\} \cup \{\pm \frac{\gamma_i - \gamma_j}{2}, 1 \leq j < i \leq r\} \cup \{\pm \frac{\gamma_i}{2}\}.$$

For this see [2] or [24]. This description yields the fact that the strongly orthogonal roots determine the domain $\omega \subset \mathfrak{a}$ in the hermitian symmetric case:

$$\omega = \{H \in \mathfrak{a} : |\gamma(H)| < \frac{\pi}{2}, \gamma \in \Gamma\}.$$

To see the inclusion $\{H \in \mathfrak{a} : |\gamma_i(H)| < \frac{\pi}{2} \forall i = 1, \dots, r\} \subset \{H \in \mathfrak{a} : |\alpha(H)| < \frac{\pi}{2} \forall \alpha \in \Psi_{\mathbb{R}}\}$, take an arbitrary $H \in \{H \in \mathfrak{a} : |\gamma_i(H)| < \frac{\pi}{2} \forall i = 1, \dots, r\}$. Then, if $\alpha = \frac{1}{2}(\pm\gamma_i \pm \gamma_j)$, $|\alpha(H)| \leq \frac{1}{2}(|\gamma_i(H)| + |\gamma_j(H)|) \leq \frac{1}{2} \cdot 2 \cdot \frac{\pi}{2} = \frac{\pi}{2}$. And if $\alpha = \pm \frac{1}{2}\gamma_i$, $|\alpha(H)| \leq \frac{1}{2}|\gamma_i(H)| \leq \frac{\pi}{4}$. The opposite inclusion is obvious since $\Gamma \subset \Sigma$.

Since $\gamma_i(X_{\gamma_j} + X_{-\gamma_j}) = 2\delta_{ij}$, one obtains the descriptions

$$\omega = \{H = \sum_{j=1}^r t_j \cdot (X_{\gamma_j} + X_{-\gamma_j}) : t_j \in (-\frac{\pi}{4}, \frac{\pi}{4})\} \quad (2)$$

and

$$\max_{\alpha \in \Sigma} |\alpha(X)| = 2 \max_{\gamma \in \Gamma} |t_\gamma|. \quad (3)$$

We now use the representation from [9] of M as a bounded symmetric domain D . The geodesic σ with $\sigma'(0) = X$ can be complexified to a proper map $\sigma : \{|Imz| < r\} \rightarrow D$, where $r = \frac{\pi}{4 \max_{\gamma \in \Gamma} |t_\gamma|}$. This follows from Corollary 7.18. In other words, the complexified geodesic exists on $\{|Imz| < 1\}$ provided that $\max_{\alpha \in \Sigma} |\alpha(X)| = \frac{\pi}{2}$.

We can now construct the required biholomorphism from the maximal Grauert domain $G \times_K (Ad(K)\omega) \subset TM$ to $M \times \overline{M}$.

First define the map as follows. Map each geodesic in $M \subset TM$ into $M \times \overline{M}$ diagonally. This map extends analytically to the complexified geodesic. The above result shows that the complexified geodesic maps into $M \times \overline{M}$ as a properly embedded disk. Doing this on each geodesic gives us our map. Considering the open embeddings of M and $\overline{M}$ in U/K and $\overline{U}/\overline{K}$ respectively, recall that the complexification $G^{\mathbb{C}} = U^{\mathbb{C}}$ acts transitively and holomorphically with parabolic isotropy groups P and $\tilde{P}$ respectively. So our map is given by

$$\begin{aligned} \Omega_{AG} &\rightarrow M \times \overline{M} \\ g \exp(iv)K^{\mathbb{C}} &\mapsto (g \exp(iv)P, g \exp(iv)\tilde{P}). \end{aligned}$$

In particular, it is holomorphic.

We check that the map is injective. If it were not injective, this would correspond to two distinct geodesics σ_1 and σ_2 in $M \hookrightarrow M \times \overline{M}$ whose complexifications intersect off the diagonal, say at a point p . Using the antiholomorphic involution, we see that the complexified geodesics actually intersect at a second point p' off the diagonal also. We note that in a Hermitian symmetric space a complexified geodesic corresponding to an initial geodesic σ with $\sigma'(0) = X$ lies inside the image of the exponential map at $\sigma(0)$ applied to the plane spanned by X and iX . Let $q_1 = \sigma_1(0)$ and $q_2 = \sigma_2(0)$. Then there is a geodesic δ in $M \times \overline{M}$ from q_1 to p and we must have $\exp_p(\delta'(p), i\delta'(p)) = \exp_{q_1}(\sigma'_1(0), i\sigma'_1(0))$ as subsets of $M \times \overline{M}$. Similarly we can find a $Y \in T_p(M \times \overline{M})$ such that $\exp_p(Y, iY) = \exp_{q_2}(\sigma'_2(0), i\sigma'_2(0))$.

If the complex subspace spanned by Y in $T_p(M \times \overline{M})$ is equal to that spanned

by $\delta'(p)$ then the original geodesics must coincide, contrary to our hypothesis. Otherwise, since the complexified geodesics also intersect at p' , we see that $\exp_p$ cannot be a diffeomorphism, which is also a contradiction. Hence, this map is injective and therefore biholomorphic onto its image.

For surjectivity, let any point $(p, q) \in M \times \overline{M}$ be given. There is a point (m, m) in the diagonal with minimal distance d to (p, q) and the length minimizing geodesic $\sigma : [0, d] \rightarrow M \times \overline{M}$ with $\sigma(0) = (m, m)$ and $\sigma(d) = (p, q)$ is orthogonal to the diagonal in (m, m) . It follows that the geodesic $\gamma : \mathbb{R} \rightarrow M \times \overline{M}$ with $\gamma(t) = \text{Exp}(J\sigma'(0))$, where J is the almost complex structure of the Kähler manifold $M \times \overline{M}$ is in fact a geodesic in the totally geodesic diagonal. Thus (p, q) is in the image of $d\gamma$ under embedding of Ω_{AG} into $M \times \overline{M}$.

Hence the map $G \times_K (Ad(K)\omega) \rightarrow M \times \overline{M}$ is a biholomorphism.

4 Proof of Theorem 7

Suppose that N is an Hermitian space of the noncompact type with an anti-holomorphic involution σ having fixed point set M . Then M is totally geodesic in N and is itself a Riemannian symmetric space with the restricted metric. In fact, if $N = G'/K'$ then $M = G/K$ where G is the centralizer of σ in G' and K is the centralizer of σ in K' .

We will now use the notation $\mathfrak{g}'$, $\mathfrak{a}'$, $\mathfrak{g}$, $\mathfrak{a}$ in the introduction for the Lie algebras and maximal abelian subalgebras of N and M respectively. We let Σ be a root system for $\mathfrak{a}$ and Σ' a root system for $\mathfrak{a}'$.

First suppose

$$\max_{\alpha \in \Sigma} |\alpha(H)| = \max_{\alpha \in \Sigma'} |\alpha(H)| \forall H \in \mathfrak{a}.$$

In this case we want to construct a biholomorphism from N to the maximal Grauert tube over M . To do this, let $\gamma \subset M \subset N$ be a geodesic. Assume that $\gamma'(0) = X \in \mathfrak{a}$ and $\max_{\alpha \in \Sigma'} |\alpha(X)| = \frac{\pi}{2}$. Then the calculation in the previous section shows that γ can be complexified in N to a proper map on

$\{|Imz| < 1\} \subset \mathbb{C}$. Now since $\max_{\alpha \in \Sigma} |\alpha(X)| = \frac{\pi}{2}$ also, identifying γ with the same geodesic in $M \subset TM$, we can extend this identification to an analytic isomorphism between the complexified geodesic in N and the differential of γ in the maximal Grauert domain.

Doing this on every geodesic, the same reasoning as in the previous section shows that we get a well-defined map from N to the maximal Grauert domain. It is a biholomorphism since it is analytic on each geodesic.

Finally, we notice that any biholomorphism from the maximal Grauert domain to N must be exactly of this form. Namely $M \subset TM$ must map to the fixed point set of an involution and, since G will push forward to a subgroup of G' , geodesics must map to geodesics. Hence the condition in Theorem 7 is both necessary and sufficient.

Remark

If the condition of Theorem 7 is not satisfied, the map constructed above from N to TM is still a biholomorphism onto its image. In this case N embeds as a subdomain of the maximal Grauert domain. Examples of this phenomenon are the Grauert domains over the symmetric spaces $SO(2,1)/SO(2)$ and $SO(3,2)/SO(3) \times SO(2)$. Here

$$SO(2,1)/SO(2) \subset SU(2,1)/S(U_2 \times U_1) \subset SO(2,1)/SO(2) \times SO(2,1)/SO(2)$$

and

$$\begin{aligned} SO(3,2)/SO(3) \times SO(2) &\subset SU(3,2)/S(U_3 \times U_2) \\ &\subset SO(3,2)/SO(3) \times SO(2) \times SO(3,2)/SO(3) \times SO(2). \end{aligned}$$

The first case here is a rank one example and so the maximal Grauert domain coincides with the maximal Grauert tube. Normalizing the curvature of the hyperbolic plane $H = SO(2,1)/SO(2)$ to be -1 , we have that $r_{max} = \frac{\pi}{2}$ and the tubes $T^{\frac{\pi}{4}}H = SU(2,1)/S(U_2 \times U_1) = B \subset \mathbb{C}^2$ and $T^{\frac{\pi}{2}}H = SO(2,1)/SO(2) \times SO(2,1)/SO(2) = D \times D \subset \mathbb{C}^2$.

5 Proof of Proposition 8

We use the same notation as in the previous section.

We first show that we can assume the Hermitian symmetric space N is irreducible. Let $N = N_1 \times \dots \times N_n$ be the decomposition of N into irreducible factors. Since the conjugation σ induces an isometry of N , it must permute the factors N_i isometrically. Hence, we can number the factors so that $\sigma(N_{2j-1}) = N_{2j}, j = 1, \dots, k$, and $\sigma(N_j) = N_j, j = 2k+1, \dots, n$. If M_j denotes the fixed point set of σ restricted to $M_{2j-1} \times M_{2j}, j = 1, \dots, k$, or $M_j, j = 2k+1, \dots, n$, it follows that M is isometric to the product of all these M_j , and that M_j is Hermitian and diagonal in $N_{2j-1} \times N_{2j}, j = 1, \dots, k$, where N_{2j-1} and N_{2j} are the same symmetric space with opposite complex structures. Note that N is a maximal Grauert domain if and only if each of the $N_j, j = 2k+1, \dots, n$ are maximal for the corresponding M_j 's, since we have already shown in section 3 that $N_{2j-1} \subset N_{2j-1} \times N_{2j}$ as above is maximal. Thus, we may assume N is irreducible.

Now suppose that $r = \text{rank}(M) = \text{rank}(N)$ but N is the maximal Grauert domain for M .

The Cartan decomposition $\mathfrak{g}' = \mathfrak{k}' + \mathfrak{p}'$ for N with respect to the Cartan involution θ yields the decomposition $\mathfrak{g} = \mathfrak{k} + \mathfrak{p}$ for M where $\mathfrak{k} = \mathfrak{k}' \cap \mathfrak{g}$ and $\mathfrak{p} = \mathfrak{p}' \cap \mathfrak{g}$.

Choose a maximal abelian subspace $\mathfrak{a} \subset \mathfrak{p}$. Without loss of generality, we can assume that $\mathfrak{a} \subset \mathfrak{a}'$, a maximal abelian subalgebra of $\mathfrak{p}'$.

Since N is irreducible, there are strongly orthogonal roots $\gamma_1, \dots, \gamma_r$ for $\mathfrak{g}'$ with respect to $\mathfrak{a}'$. In the root decomposition

$$\mathfrak{g}' = Z(\mathfrak{a}') + \sum_{\alpha \in \Sigma'} \mathfrak{g}'^{\alpha} \quad (4)$$

it is known that $\dim(\mathfrak{g}'^{\gamma_i}) = 1$ for all i , see for instance [2], section 2.3.

Now, associated to γ_i there is a subalgebra $\mathfrak{sl}_2(\mathbb{R}) \subset \mathfrak{g}'$ spanned by $\mathfrak{g}'^{\gamma_i}$,

$\mathfrak{g}'^{-\gamma_i} = \theta(\mathfrak{g}'^{\gamma_i})$ and some $x_i \in \mathfrak{a}'$, see again [2]. Upon exponentiation, this $\mathfrak{sl}_2(\mathbb{R})$ generates a complex disk in N .

The root space decomposition of Hermitian symmetric spaces discussed in section 3 implies that

$$|\gamma_i(x_i)| > |\alpha(x_i)| \forall \alpha \in \Sigma' \setminus \{\pm\gamma_i\}. \quad (5)$$

If the maximality condition holds, then there exists a $\beta \in \Sigma$ such that $\beta(x_i) = \gamma_i(x_i)$ and hence $\beta_i = \gamma_i$. Since $\dim(\mathfrak{g}'^{\gamma_i}) = 1$, we have that $\mathfrak{g}'^{\gamma_i} \subset \mathfrak{g}$ and also $\mathfrak{g}'^{-\gamma_i} \subset \mathfrak{g}$. Therefore the $\mathfrak{sl}_2(\mathbb{R})$ associated to γ_i is a subalgebra of $\mathfrak{g}$. This contradicts the fact that M is a totally-real submanifold of N .

Now suppose that $s = \text{rank}(N) > \text{rank}(M) = r$. Let J be the complex structure on $\mathfrak{p}' = T_0(N)$. Then $\mathfrak{a}' = \mathfrak{a} + \mathfrak{a}''$ where $\mathfrak{a}''$ is contained in the (-1) -eigenspace of the complex conjugation σ on N and $J\mathfrak{a}'' \subset \mathfrak{p}$. Here again $\mathfrak{a}'$ is a maximal abelian subspace of $\mathfrak{p}'$ and $\mathfrak{a}$ is maximal abelian in $\mathfrak{p}$.

The polydisk theorem (see for instance [2] again) tells us that $\exp(\mathfrak{a}')$ can be complexified to a totally geodesic, embedded polydisk $\Delta^s \subset N$. The involution σ restricts to an involution on Δ^s whose fixed-point set P is totally geodesic in Δ^s and hence M . Clearly $T_0(P) = \mathfrak{a} + J\mathfrak{a}''$ and P has rank r and dimension s , since $\mathfrak{a} \cap J\mathfrak{a}'' = 0$: otherwise the bounded symmetric domain would contain a J -invariant complete 2-dimensional flat, which would be a copy of $\mathbb{C}$, contradicting Liouville's theorem.

Now, an involution on a polydisk Δ^s either preserves a factor or preserves a pair of factors Δ^2 , permuting the pair. In the second case, up to choosing coordinates $(z, w) \in \Delta^2$ we may take $\sigma(z, w) = (\bar{w}, \bar{z})$ and the fixed-point set is a copy of the hyperbolic plane H . Thus $P = H^p \times \mathbb{R}^q$. The $\mathbb{R}$ -factors come from factors of Δ^s fixed by σ and so we may assume that they are tangent to some $x_i \in \mathfrak{a}$, as above, corresponding to a strongly orthogonal root. Comparing ranks and dimensions we find that $p + q = r$ and $2p + q = s$ respectively. Thus $p = s - r$ and $q = 2s - r$.

Suppose that $s < 2r$. Then P contains at least one $\mathbb{R}$ -factor and so we may assume that there exists an $x_i \in T_0(P) \subset T_0(M) = \mathfrak{p}$. If N were the maximal Grauert domain then we could find an $X \in \mathfrak{g}$ such that $[X, x_i] = \gamma_i(x_i)X$. We claim that $X \in \mathfrak{g}'^{\gamma_i}$. But by the direct sum decomposition (4) above we have

$$X = z + \sum_{\alpha \in \Sigma'} Y_\alpha$$

where $Y_\alpha \in \mathfrak{g}'^\alpha$. Hence

$$\gamma_i(x_i)z + \sum_{\alpha \in \Sigma'} (\gamma_i(x_i) - \alpha(x_i))Y_\alpha = 0.$$

By the inequality (5) this gives us that $z = 0$ and $Y_\alpha = 0$ for $\alpha \neq \gamma_i$, justifying our claim. Now we can argue as above to produce a holomorphic disk in M and derive a contradiction as before.

Finally suppose that $s = 2r$. (Certainly $s \leq 2r$ as $J\mathfrak{a}''$ is abelian in $\mathfrak{p}$, and so is of dimension $\leq r$.) In this case we would like to show that N is the maximal Grauert domain. To do this, given a vector $X \in \mathfrak{p}$, we need to find a Jacobi field along the corresponding geodesic which when complexified has a pole at the boundary of the complexified geodesic (see section 2). We may assume that $X \in \mathfrak{a}$ and clearly it would suffice to find such a Jacobi field tangent to P . Since $s = 2r$, by the argument in the preceding paragraph, we have that $P = H^r$. Each copy of H in this factorization corresponds to two strongly orthogonal roots, say, γ_1, γ_2 permuted up to sign by σ . The diagonal copy of $Sl_2(\mathbf{R})$ in the product of the $Sl_2(\mathbf{R})$'s corresponding to γ_1, γ_2 acts transitively on H , and the product group acts transitively on the corresponding bidisk Δ^2 in our maximal Δ^s . As each such H is Hermitian, from section 3 we know that the corresponding bidisk Δ^2 is its maximal Grauert domain, as Δ^s is of $P = H^r$. Finally, since P is totally geodesic both in M and in Δ^s which in turn are totally geodesic in N , a Jacobi field (in P or in M , equivalently) does exist tangent to P as required.

6 Proof of Theorem 6

Let $M = G/K$ be an irreducible Riemannian symmetric space of noncompact type and N an associated Grauert domain. It is the aim of this section to show that N is either rigid or Hermitian symmetric.

We will use the fact that as N admits a bounded strictly plurisubharmonic function, namely the length-squared function ρ^2 , it is a hyperbolic complex manifold, see [20], Theorem 3. Let d denote the metric space structure on N given by integrating the infinitesimal Kobayashi metric.

First note that any biholomorphism of N which preserves M must be the differential of an isometry of M . Such a result was proven in [15], section 6, in the context of Grauert tubes of less than maximal radius. The proof, however, extends to our more general case. We remark here that the tubes of less than maximal radius are all complete hyperbolic. We do not know how to prove this for general Grauert domains (except as a consequence of some of them being Hermitian symmetric). The proof of [15] as written does use tautness of the tubes, but in fact hyperbolicity is enough, see [16], Chapter V, Theorem 3.3.

Suppose that N is not rigid. Then we can say that there exists a biholomorphism of N which moves M off itself. According to [16], Chapter V, Theorem 2.1, the biholomorphism group $Aut_{\mathbb{C}}(N)$ is a Lie group with compact isotropy groups. As the orbit of a point in $N \setminus M$ under the action of G has higher dimension than M , we deduce that the identity component $Aut_{\mathbb{C}}^0(N)$ must itself move M off itself. We think of elements of the Lie algebra of $Aut_{\mathbb{C}}^0(N)$ as vector fields on N . Then given a base point $p \in M$, there exists a vector field in the Lie algebra transverse to M at p . We now use a result of Cartan and the fact that M is irreducible to find that in fact vector fields in the Lie algebra must span $T_p(TM)$, because of course the Lie algebra is invariant under the action of $Ad(K)$.

As a consequence of this, there exist elements of $Aut_{\mathbb{C}}^0(N)$ taking p to all points in a sufficiently small neighbourhood U of p in N . We claim this implies

that (N, d) must be complete.

Proof of claim

Again we argue by contradiction. If (N, d) were not complete, there exists a non-empty collection $\mathcal{C}$ of Cauchy sequences $\{x_i\}$ with no convergent subsequences.

Hence the function $f : N \rightarrow [0, \infty)$ given by

$$f(q) = \inf_{\mathcal{C}} \sup_i d(q, x_i)$$

is in fact finite-valued.

As d is invariant under $\text{Aut}_{\mathbb{C}}(N)$, so is f and hence f is constant, say equal to D , in U . Choose $\epsilon > 0$ such that $S = \{q \in N \mid d(p, q) = \epsilon\}$ is a compact subset of U .

Now let $\{x_i\} \in \mathcal{C}$ be such that $\sup_i d(p, x_i) < D + \frac{\epsilon}{2}$.

Every path from p to x_i must pass through a point in S . Therefore we can find a sequence $q_i \in S$ with $d(q_i, x_i) < D - \frac{\epsilon}{2}$.

As S is compact, after taking subsequences we may assume that q_i converges to a point $q \in S$.

Then replacing the subsequence $\{x_i\}$ by $\{x_i\}_{i \geq N}$ for some large N , we have

$$\sup_i d(q, x_i) \leq \sup_i (d(q, q_i) + d(q_i, x_i)) < D - \frac{\epsilon}{4}.$$

But this contradicts $f(q) = D$, completing the proof of the claim.

We already know that the set of points in N which can be mapped to p by an element of $\text{Aut}_{\mathbb{C}}(N)$ is open, but now since N is complete hyperbolic and in particular taut, this set is also closed. Hence $\text{Aut}_{\mathbb{C}}(N)$ acts transitively. Also, the geodesic symmetry at p of M extends to an involution in $\text{Aut}_{\mathbb{C}}(N)$, and the fixed point set of the action by conjugation is exactly the (compact) isotropy group of p . Thus N is an Hermitian symmetric space as required.

7 Direct Proof that the Akhiezer-Gindikin Domain is Stein.

Theorem 10 *Let $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$ be the Cartan decomposition, $\mathfrak{a} \subset \mathfrak{p}$ maximal abelian, W the Weyl group for the adjoint action of K on $\mathfrak{p}$ and $\omega \subset \mathfrak{a}$ the set describing the Akhiezer-Gindikin domain*

$$\Omega_{AG} = \{g \exp(i\xi) K^{\mathbb{C}}, g \in G, \xi \in \omega\}.$$

Let u be a strictly convex, W -invariant, smooth function on ω and $\tilde{u}$ its G -invariant extension on Ω_{AG} . Then $\tilde{u}$ is strictly plurisubharmonic.

Proof

Let us set the notation. The symbols $\mathfrak{g}, \mathfrak{k}, \mathfrak{p}, \mathfrak{a}, \theta$ denote the Lie algebras, Cartan involution, eigenspaces, etc. as above, associated to our symmetric space M . Let $\mathfrak{g}^{\mathbb{C}} = \mathfrak{g} \otimes \mathbb{C}, \mathfrak{k}^{\mathbb{C}}, \theta$, etc., be the corresponding complexified objects. Let $G, K, G^{\mathbb{C}}, K^{\mathbb{C}}$, etc., be the corresponding groups and complex groups. Let W be the Weyl group of $\mathfrak{g}$ with respect to $K, \mathfrak{a}$. Let $X = G^{\mathbb{C}}/K^{\mathbb{C}}$ be the affine complexification of M . Let $\mathfrak{u}$ be the compact twin of $\mathfrak{g}$, where $\mathfrak{u} = \mathfrak{k} \oplus i\mathfrak{p}$, where multiplication by i is meant in $\mathfrak{g}^{\mathbb{C}}$, and let $U \subset G^{\mathbb{C}}$ be the corresponding subgroup, the maximal compact subgroup of $G^{\mathbb{C}}$. Let $\xi_1, \xi_2, \dots, \xi_n$ be an orthonormal basis of $\mathfrak{p}$. There is an orthogonal direct sum decomposition $\mathfrak{p} = \mathfrak{a} + \mathfrak{p}'$ where $\mathfrak{p}'$ is the orthogonal complement of $\mathfrak{a} \subset \mathfrak{p}$. Let $M^c = U/K \subset X$ be the compact twin symmetric space of our original M . If σ denotes the conjugation of X fixing M , let $\tau = \theta \circ \sigma = \sigma \circ \theta$ be the conjugation for M^c . Let Ω_{AG} be the Akhiezer-Gindikin domain, which is the image $G \cdot \exp(iAd(K)\omega) \cdot K^{\mathbb{C}}$ in $G^{\mathbb{C}}/K^{\mathbb{C}}$. We write here Ω_{AG} for $G \times_K Ad(K)\omega$, too. Here the $\exp$ can be taken either as group exponential in $G^{\mathbb{C}}$, or as the geodesic exponential in the compact twin M^c . For any vector $\xi \in \mathfrak{g} \otimes \mathbb{C}$, let $\tilde{\xi}$, or sometimes $[\xi]$, denote the corresponding vectorfield given by $\tilde{\xi}(p) = \frac{d}{dt} \exp(t\xi) \cdot p \big|_{t=0}$ on either X , or equivalently on $Ad(K)\omega \subset i\mathfrak{p}$, for fields tangent to M^c within Ω_{AG} . Let u be a

smooth, strictly convex, W -invariant function on $\mathfrak{a}$, and let $\tilde{u}$ be the corresponding function on Ω_{AG} . We wish to show that $\tilde{u}$ is strictly plurisubharmonic. Fix $\xi_0 \in \text{Ad}(K)\omega$; we want to show $i\partial\bar{\partial}(\tilde{u}) > 0$ at $x_0 = \exp(i\xi_0) \cdot K \in M^c$. Without loss of generality, we can assume $\xi_0 \in \omega$, in particular, $\xi_o \in \mathfrak{a}$.

The only way we use that we are inside Ω_{AG} at x_0 is in the following observation: $\tilde{\xi}_1, \dots, \tilde{\xi}_n$ are linearly independent at x_0 , and span a totally real subspace of the tangent space of X at x_0 . This follows immediately from the (equivariant) bundle structure of the Grauert domain construction.

Since the group $G^{\mathbb{C}}$ acts holomorphically on X , we have that $Z_k = \tilde{\xi}_k^{1,0}$ is a holomorphic vector field, and $\bar{Z}_j$ is anti-holomorphic. As a result, $[Z_l, \bar{Z}_k] = 0$, for all $k, l = 1, \dots, n$. Since the span of the $\tilde{\xi}_k$ is totally real, the complex fields $Z_1, \dots, Z_n$ are $\mathbb{C}$ -linearly independent at x_0 , and span the complex tangent space at that point. Thus, it suffices to show the matrix $i\partial\bar{\partial}(\tilde{u})(Z_i, \bar{Z}_j)$ is positive definite. This reduces immediately to the matrix $Z_i \cdot \bar{Z}_j(\tilde{u})$.

Write Z_k as $\frac{1}{2}(\tilde{\xi}_k - iJ\tilde{\xi}_k)$, and therefore,

$$\begin{aligned} Z_k \bar{Z}_l(\tilde{u}) &= \frac{1}{4}(\tilde{\xi}_k - iJ\tilde{\xi}_k)(\tilde{\xi}_l + iJ\tilde{\xi}_l)(\tilde{u}) \\ &= \frac{1}{4}(i\tilde{\xi}_k J\tilde{\xi}_l + J\tilde{\xi}_k J\tilde{\xi}_l)(\tilde{u}), \end{aligned}$$

since $\tilde{\xi}_l(\tilde{u}) \equiv 0$, for all l . Note also that $J\tilde{\xi}_k = \widetilde{(i\xi_k)}$, for all k . Finally, note that

$$\tilde{\xi}_k J\tilde{\xi}_l(\tilde{u}) = [\tilde{\xi}_k, J\tilde{\xi}_l](\tilde{u}) = (i[\xi_k, \xi_l])^\sim(\tilde{u}) = J([\xi_k, \xi_l])^\sim(\tilde{u}),$$

since the flow of $\tilde{\xi}_k$ is holomorphic. Now $[\xi_k, \xi_l] = \eta \in \mathfrak{k}$, and as $\mathfrak{k}$ acts on M^c at x_0 , we have the basic observation that

the tangent space to the orbit of K through x_0 is spanned by vectors $\tilde{i\xi}, \xi \in \mathfrak{p}'$.

(Notice that these two subspaces of tangent vectors are equal at points $x_0 = \exp(i\xi_0) \cdot 0$, where $\xi_0 \in \mathfrak{p}$ is a regular element, and hence the claim holds for any

$\xi_0 \in \mathfrak{p}$.) Hence, at x_0 there is some $\xi \in \mathfrak{p}$ such that

$$(i[\xi_k, \xi_l])^\sim = J([\xi_k, \xi_l])^\sim = Ji\xi = -\xi$$

at x_0 . But then this implies that

$$\tilde{\xi}_k J \tilde{\xi}_l(\tilde{u}) = [\tilde{\xi}_k, J \tilde{\xi}_l](\tilde{u}) = -\tilde{\xi}(\tilde{u}) = 0,$$

at x_0 , again, by G -invariance of $\tilde{u}$. So, we have reduced our task to showing that the matrix

$$\widetilde{i\xi_k i\xi_l}(\tilde{u})$$

is positive definite at x_0 .

We now write the basis $\xi_1, \xi_2, \dots, \xi_n$ of $\mathfrak{p}$ so that $\xi_i, \xi_j, j, i = 1, \dots, r = \dim(\mathfrak{a})$, denote elements in $\mathfrak{a}$, and ξ_k, ξ_l from among $\xi_{r+1}, \dots, \xi_n$ are in $\mathfrak{p}' =$ orthogonal complement of $\mathfrak{a} \subset \mathfrak{p}$. More precisely, let Σ be the set of roots for $\mathfrak{a}$ acting on $\mathfrak{g}$, and let Σ_+ be the positive roots with respect to some ordering. For $\alpha \in \Sigma$, let X_α be a non-zero vector in $\mathfrak{g}^\alpha = \{Y \in \mathfrak{g} \mid [H, Y] = \alpha(H)Y, \text{ for all } H \in \mathfrak{a}\}$. For $\alpha \in \Sigma$, $X_\alpha - \theta X_\alpha \in \mathfrak{p}'$, and in fact, a basis for $\mathfrak{p}'$ is given by $X_\alpha - \theta X_\alpha, \alpha \in \Sigma_+$. Note that in this notation we are counting the α 's $\in \Sigma_+$ with *multiplicity*. Finally, the $X_\alpha - \theta X_\alpha$ are orthogonal to one another, and so, up to scale, we may take them as an orthonormal basis for $\mathfrak{p}'$. In other words, for $k, l = r+1, \dots, n$, we can take ξ_k, ξ_l to be of the form $X_\alpha - \theta X_\alpha$, up to a scale factor. Let $x_1, \dots, x_n$ denote the affine coordinates in $\mathfrak{p}$ corresponding to this (orthonormal) basis of $\mathfrak{p}$.

We have three types of terms to consider in the matrix $\widetilde{i\xi_s i\xi_t}(\tilde{u})$:

- i) $\mathfrak{a}$ -terms $\widetilde{i\xi_i i\xi_j}(\tilde{u}), i, j = 1, \dots, r$.
- ii) cross-terms $\widetilde{i\xi_i i\xi_k}(\tilde{u}), i = 1, \dots, r, k = r+1, \dots, n$.
- iii) $\mathfrak{p}'$ -terms $\widetilde{i\xi_k i\xi_l}(\tilde{u}), k, l = r+1, \dots, n$.

In case ii), notice that $[Z_s, \bar{Z}_t] \equiv 0$ and the calculation above imply that $\widetilde{i\xi_i, i\xi_k}(\tilde{u}) \equiv 0$ also.

The first two types of terms are easy to compute, and do not even require us to know the fields $\widetilde{i\xi_s}$ very explicitly.

Case i): In this case, the subalgebra $\mathfrak{a}$ is abelian, and the subspace $\exp(i\mathfrak{a}) \cdot 0 \subset M^c$ is flat. Therefore,

$$\widetilde{i\xi_i}\widetilde{i\xi_j}(\tilde{u})(x_0) = \frac{\partial^2 u}{\partial x_i \partial x_j}(\xi_0).$$

Case ii): First use the basic observation above to see that, at all regular $\xi_0 \in \mathfrak{a}$, the tangent vector $\widetilde{i\xi_k}$ is tangent to the orbit of K through $x_0 = \exp(i\xi_0) \cdot 0$, and by K -invariance of $\tilde{u}$, $\widetilde{i\xi_k}(\tilde{u})(x_0) = 0$. Since the regular ξ_0 are dense in $\mathfrak{a}$, this is true at every $x_0 \in \exp(i\mathfrak{a}) \cdot 0$. Therefore, since every $\widetilde{i\xi_i}$ is tangent to $\exp(i\mathfrak{a}) \cdot 0$, we have that $\widetilde{i\xi_i}\widetilde{i\xi_k}(\tilde{u})(x_0) = 0$.

Case iii): Here we have to examine the differential of the exponential map more clearly to see how the fields $\widetilde{i\xi_s}$ are transported from M^c back to $\mathfrak{p}$ by the inverse of the exponential map.

We will work along constant speed geodesics $\gamma_\xi(t) = \exp(it\xi) \cdot 0, \xi \in \mathfrak{p}$, in M^c . Since we will be using the Jacobi equation, it is useful to understand parallel transport along γ_{ξ_0} explicitly. Let $g_t = \exp(it\xi_0) \in U$, note that the tangent space to M^c at $g_t = \gamma_{\xi_0}(t)$ is identified with $dg_{t*}\mathfrak{p} = \mathfrak{u} \bmod \text{Ad}(g_t)(\mathfrak{k})$. Parallel vectors along γ_{ξ_0} are simply those of the form $dg_{t*}(\xi)$ for fixed $\xi \in \mathfrak{p}$. Given this identification, a Jacobi field $Y(t)$ can be written in terms of the parallel fields as

$$Y(t) = dg_{t*}\left(\sum_{j=1}^r v_j(t)i\xi_j + \sum_{\alpha \in \Sigma_+} v_\alpha(t)i(X_\alpha - \theta X_\alpha)\right),$$

and the coefficient functions satisfy the ordinary differential equations:

$$\ddot{v}_j \equiv 0,$$

$$\ddot{v}_\alpha = \{\alpha(i\xi_0)\}^2 v_\alpha = -\{\alpha(\xi_0)\}^2 v_\alpha.$$

Taking into account the initial conditions $v_\alpha(0) = 0, \dot{v}_\alpha(0) = 1$, and setting all other coefficients equal to 0, we get that

$$d \exp_*(i\xi_0) : i(X_\alpha - \theta X_\alpha) \longrightarrow dg_{1*} \left(\frac{\sin(\alpha(\xi_0))}{\alpha(\xi_0)} i(X_\alpha - \theta X_\alpha) \right).$$

We are assuming, provisionally, that ξ_0 is regular, so $\alpha(\xi_0) \neq 0$, for all $\alpha \in \Sigma$. If we denote the inverse map to $\exp$ by “ $\log$ ”, then we have

$$d \log_*(x_0) : dg_{1*}(i(X_\alpha - \theta X_\alpha)) \longrightarrow \frac{\alpha(\xi_0)}{\sin(\alpha(\xi_0))} i(X_\alpha - \theta X_\alpha).$$

Recall that, since $0 \neq \xi_0 \in \omega_0$, we have $0 < |\alpha(\xi_0)| < \frac{\pi}{2}$, so the denominator above doesn't vanish.

Next we have to figure out how to represent the fields $\widetilde{i\xi_k}$ back on $\mathfrak{p}$. Of course, these are just the derivatives of the actions of $\exp(it(X_\alpha - \theta X_\alpha)) \in U$ on M^c . This is computable at $x_0 = g_1$ as $dg_{1*}[\frac{d}{dt}|_{t=0} g_1^{-1} \cdot \exp(it(X_\alpha - \theta X_\alpha)) \cdot g_1 \cdot 0]$. Now this curve at $0 \in M^c$ obviously has as derivative at $t = 0$ the image of $Ad(g_1^{-1})i(X_\alpha - \theta X_\alpha) \in \mathfrak{u}$ modulo $\mathfrak{k}$. We can calculate this using $g_1 = \exp(i\xi_0) \in U$. So,

$$\begin{aligned} Ad(g_1^{-1})i(X_\alpha - \theta X_\alpha) &= ie^{-iad(\xi_0)}(X_\alpha - \theta X_\alpha) \\ &= e^{-i\alpha(\xi_0)}iX_\alpha - ie^{i\alpha(\xi_0)}\theta X_\alpha \\ &= \cos(\alpha(\xi_0))i(X_\alpha - \theta X_\alpha) - \sin(\alpha(\xi_0))(X_\alpha + \theta X_\alpha) \\ &= \cos(\alpha(\xi_0))i(X_\alpha - \theta X_\alpha) \bmod \mathfrak{k}, \end{aligned}$$

since $X_\alpha + \theta X_\alpha \in \mathfrak{k}$, for every $\alpha \in \Sigma$. From this we conclude

$$d \log_*(x_0) : [i(X_\alpha - \theta X_\alpha)]^\sim \longrightarrow \frac{\alpha(\xi_0) \cos(\alpha(\xi_0))}{\sin(\alpha(\xi_0))} i(X_\alpha - \theta X_\alpha).$$

Now, before computing $\widetilde{i\xi_k} \widetilde{i\xi_l}(\tilde{u})(x_0)$, we make an auxiliary computation of $\widetilde{i\xi_k} \widetilde{i\xi_l}(\tilde{u}_0)(x_0)$, where u_0 denotes the gradient of u on $\mathfrak{p}$ at ξ_0 , and by abuse of notation, will also denote the function $\xi \rightarrow u_0 \cdot \xi$ determined by taking the inner product of ξ with u_0 . Similarly, $\tilde{u}_0$ denotes the same function transported

to $\exp(i\text{Ad}(K)\omega_0) \cdot 0 \subset M^c$ by the exponential function. We carry out the computation on $\mathfrak{p}$ near ξ_0 , that is, we compute

$$\begin{aligned} & [i(X_\alpha - \theta X_\alpha)]^\sim [i(X_\beta - \theta X_\beta)]^\sim (\tilde{u}_0)(x_0) \\ &= d\log_*([i(X_\alpha - \theta X_\alpha)]^\sim(x_0))(u_0 \cdot d\log_*([i(X_\beta - \theta X_\beta)]^\sim)), \end{aligned}$$

evaluated at $\xi_0 \in \mathfrak{p}$.

We first want to use the basic observation above to replace the tangent vector $[i(X_\alpha - \theta X_\alpha)]^\sim$ at x_0 by $\tilde{\eta}$, for suitable $\eta \in \mathfrak{k}$. We repeat that we are assuming provisionally that ξ_0 is regular. To do this, let us calculate much as before

$$\begin{aligned} (X_\alpha + \theta X_\alpha)^\sim(x_0) &= dg_{1*}[\frac{d}{dt}|_{t=0} g_1^{-1} \cdot \exp(t(X_\alpha + \theta X_\alpha)) \cdot g_1 \cdot 0] \\ &= dg_{1*}(e^{-i \text{ad}(\xi_0)}(X_\alpha + \theta X_\alpha)) \\ &= dg_{1*}(\cos(\alpha(\xi_0))(X_\alpha + \theta X_\alpha) - i \sin(\alpha(\xi_0))(X_\alpha - \theta X_\alpha)) \\ &= dg_{1*}(-\sin(\alpha(\xi_0))i(X_\alpha - \theta X_\alpha)) \text{mod } \mathfrak{k}. \end{aligned}$$

Taken together with what was shown above, we conclude

$$[i(X_\alpha - \theta X_\alpha)]^\sim(x_0) = -\cot(\alpha(\xi_0))(X_\alpha + \theta X_\alpha)^\sim.$$

Going back to our original computation, this gives us that

$$\begin{aligned} & [i(X_\alpha - \theta X_\alpha)]^\sim [i(X_\beta - \theta X_\beta)]^\sim (\tilde{u}_0)(x_0) \\ &= -\cot(\alpha(\xi_0))d\log_*([i(X_\alpha + \theta X_\alpha)]^\sim(x_0))(u_0 \cdot d\log_*([i(X_\beta - \theta X_\beta)]^\sim)). \end{aligned}$$

Because the subgroup K fixes $0 \in M^c$, the exponential map from 0 is equivariant with respect to the action of K on $T_0(M^c) = \mathfrak{p}$ and on M^c . Thus, we conclude that

$$\begin{aligned} & [i(X_\alpha - \theta X_\alpha)]^\sim [i(X_\beta - \theta X_\beta)]^\sim (\tilde{u}_0)(x_0) \\ &= -\cot(\alpha(\xi_0))\frac{d}{dt}[u_0 \cdot d\log_*([i(X_\beta - \theta X_\beta)]^\sim)(\text{Ad}(\exp(t(X_\alpha + \theta X_\alpha)))\xi_0)]|_{t=0}. \end{aligned}$$

For simplicity, set $k_t = \exp(t(X_\alpha + \theta X_\alpha)) \in U$. K -equivariance of $\exp$ implies

$$\begin{aligned}
& \frac{d}{dt} [u_0 \cdot d \log_* ([i(X_\beta - \theta X_\beta)]^\sim)(Ad(k_t)\xi_0)]|_{t=0} \\
&= u_0 \cdot \left[\frac{d}{dt} (Ad(k_t) \cdot d \log_*(x_0) \{ dk_t^{-1} ([i(X_\beta - \theta X_\beta)]^\sim) \}) \right]_{t=0} \\
&= u_0 \cdot \left[\frac{d}{dt} (Ad(k_t) \cdot d \log_*(x_0) \{ [iAd(k_t^{-1})(X_\beta - \theta X_\beta)]^\sim \}) \right]_{t=0} \\
&= u_0 \cdot [X_\alpha + \theta X_\alpha, d \log_*(x_0) ([i(X_\beta - \theta X_\beta)]^\sim)] \\
&\quad - u_0 \cdot d \log_*(x_0) ([i[X_\alpha + \theta X_\alpha, X_\beta - \theta X_\beta]]^\sim).
\end{aligned}$$

We next compute $[X_\alpha + \theta X_\alpha, i(X_\beta - \theta X_\beta)] = i([\theta X_\alpha, X_\beta] - \theta[\theta X_\alpha, X_\beta])$. Notice that if $\alpha \neq \beta$, then $i([\theta X_\alpha, X_\beta] - \theta[\theta X_\alpha, X_\beta]) \in i\mathfrak{p}'$, which is tangent to the K -orbit of ξ_0 . Hence $u_0 \cdot i([\theta X_\alpha, X_\beta] - \theta[\theta X_\alpha, X_\beta]) = 0$.

For the second term above, we note that $d \log_*(x_0)$ sends the subspace spanned by $[i(X_\alpha - \theta X_\alpha)]^\sim$ to the subspace spanned by $i(X_\alpha - \theta X_\alpha)$, and similarly the subspace spanned by $i\widetilde{\xi}_j, j = 1, \dots, r$, gets sent to the subspace $i\mathfrak{a} \subset i\mathfrak{p}$. As a result,

$$\begin{aligned}
& u_0 \cdot d \log_*(x_0) (-[i[X_\alpha + \theta X_\alpha, X_\beta - \theta X_\beta]]^\sim) \\
&= u_0 \cdot d \log_*(x_0) ([i\{[\theta X_\alpha, X_\beta] - \theta[\theta X_\alpha, X_\beta] + [X_\alpha, X_\beta] - \theta([X_\alpha, X_\beta])\}]^\sim) = 0,
\end{aligned}$$

for all $\alpha \neq \beta \in \Sigma_+$.

Finally, in the case when $\alpha = \beta$, then $i([\theta X_\alpha, X_\alpha] - \theta[\theta X_\alpha, X_\alpha]) \in iZ(\mathfrak{a}) \cap \mathfrak{a} = i\mathfrak{a}$. Recalling that $d \log_*(x_0)$ is the “identity” when restricted to vectors tangent to the flat $\exp(i\mathfrak{a}) \cdot 0 \subset M^c$, we get

$$u_0 \cdot d \log_*(x_0) ([i\{[\theta X_\alpha, X_\alpha] - \theta[\theta X_\alpha, X_\alpha]\}]^\sim) = u_0 \cdot i\{[\theta X_\alpha, X_\alpha] - \theta[\theta X_\alpha, X_\alpha]\},$$

where $i\{[\theta X_\alpha, X_\alpha] - \theta[\theta X_\alpha, X_\alpha]\} \in \mathfrak{a}$. Putting this together with what we have left of the first term, we get

$$[i(X_\alpha - \theta X_\alpha)]^\sim [i(X_\beta - \theta X_\beta)]^\sim (\tilde{u}_0)(x_0) = 0, \text{ if } \alpha \neq \beta,$$

and

$$\begin{aligned} & [i(X_\alpha - \theta X_\alpha)]^\top [i(X_\alpha - \theta X_\alpha)]^\top (\tilde{u}_0)(x_0) \\ &= u_0 \cdot i([\theta X_\alpha, X_\alpha] - \theta[\theta X_\alpha, X_\alpha]) - u_0 \cdot i([\theta X_\alpha, X_\alpha] - \theta[\theta X_\alpha, X_\alpha]) = 0, \end{aligned}$$

for all $\alpha \in \Sigma_+$.

Summarizing, we have shown

$$[i(X_\alpha - \theta X_\alpha)]^\top [i(X_\beta - \theta X_\beta)]^\top (\tilde{u}_0)(x_0) = 0, \text{ for all } \alpha, \beta \in \Sigma_+.$$

By our choice of u_0 as the gradient of u at ξ_0 , we have that the function $u - u_0$ has a critical point at $\xi_0 \in \mathfrak{p}$, equivalently, $\tilde{u} - \tilde{u}_0$ has a critical point at x_0 . (Recall that, by abuse of notation, we are denoting by u_0 the linear function $u_0 \cdot \xi$.) Hence the derivatives

$$[i(X_\alpha - \theta X_\alpha)]^\top [i(X_\beta - \theta X_\beta)]^\top (\tilde{u})(x_0) = [i(X_\alpha - \theta X_\alpha)]^\top [i(X_\beta - \theta X_\beta)]^\top (\tilde{u} - \tilde{u}_0)(x_0)$$

can be evaluated in terms of the invariantly defined Hessian of $u - u_0$ at $\xi_0 \in \mathfrak{p}$, that is, we get

$$\begin{aligned} & [i(X_\alpha - \theta X_\alpha)]^\top [i(X_\beta - \theta X_\beta)]^\top (\tilde{u} - \tilde{u}_0)(x_0) \\ &= \text{Hess}(u - u_0)(\xi_0) (d \log_*(x_0)([i(X_\alpha - \theta X_\alpha)]^\top), d \log_*(x_0)([i(X_\beta - \theta X_\beta)]^\top)) \\ &= \alpha(\xi_0) \cot(\alpha(\xi_0)) \beta(\xi_0) \cot(\beta(\xi_0)) \text{Hess}(u - u_0)(\xi_0)(i(X_\alpha - \theta X_\alpha), i(X_\beta - \theta X_\beta)). \end{aligned}$$

This last matrix is obviously positive definite when ξ_0 is regular, since u is strictly convex. For ξ_0 not regular, the result follows from the regular case, passing to the limit ξ_0 from regular $\xi \in \mathfrak{a}$, taking into account that the function

$$\alpha(\xi) \cot(\alpha(\xi)) = \frac{\alpha(\xi)}{\sin(\alpha(\xi))} \cdot \cos(\alpha(\xi))$$

has a finite, non-vanishing limit at ξ_0 as long as $|\alpha(\xi_0)| < \frac{\pi}{2}$. This completes the proof of the theorem.

Corollary 11 *The domain Ω_{AG} has the Stein property.*

Proof

Let Γ be a discrete subgroup of G acting completely discontinuously on $M = G/K$ such that $\Gamma \backslash M$ is a compact manifold. Such a Γ can be found according to [4]. Let $\tilde{u}$ be given as in the theorem by a strictly convex, W -invariant function u compactly exhausting ω . For instance,

$$u(\xi) := \sum_{\alpha \in \Sigma} \frac{1}{(\frac{\pi}{2})^2 - \alpha(\xi)^2}$$

has these properties with respect to the symmetric space metric on $\mathfrak{p} \cong T_{eK}G/K$ restricted to $\mathfrak{a}$. Then $\tilde{u}$ pushes down to a smooth, strictly plurisubharmonic exhaustion function of $\Gamma \backslash \Omega_{AG}$, which is therefore Stein. The universal covering Ω_{AG} of $\Gamma \backslash \Omega_{AG}$ is then also Stein (see [21], p. 66 f.) completing the proof of the corollary.

References

- [1] D. N. Akhiezer and S. G. Gindikin, *On Stein extensions of real symmetric spaces*, Math. Ann., 286(1990), no.1-3, 1-12.
- [2] A. Ash, D. Mumford, M. Rapoport, Y. Tai, *Smooth compactification of locally symmetric varieties*, Lie Groups: History, Frontiers and Applications, Vol. IV, Math. Sci. Press, Brookline, Mass., 1975.
- [3] H. Azad and J.J. Loeb, *Plurisubharmonic functions and Kählerian metrics on complexification of symmetric spaces*, Indag. Math., 3(1990), no. 4, 365-375.
- [4] A. Borel, *Compact Clifford-Klein forms of Symmetric Spaces*, Topology, Vol. 2 (1964), pp. 111-122
- [5] D. Burns, *On the uniqueness and characterization of Grauert tubes*, Complex analysis and geometry (Trento, 1993), 119-133, Lecture Notes in Pure and Appl. Math. 173.

- [6] D. Burns and R. Hind, *Symplectic Geometry and the Uniqueness of Grauert tubes*, J. Geom. and Funct. Anal., 11 (2001), 1-10.
- [7] V. Guillemin and M. Stenzel, *Grauert tubes and the homogeneous Monge-Ampère equation*, J. Diff. Geom., 34 (1991), 561-570.
- [8] S. Halverscheid, *Maximal Domains of Definition of Adapted Complex Structures for Symmetric Spaces of Non-compact Type*, Thesis, Ruhr-Universität Bochum, 2001
- [9] S. Helgason, *Differential Geometry, Lie Groups and Symmetric Spaces*, Pure and Applied Math., 80, Academic Press, New York-London, 1978.
- [10] H. A. Jaffee, *Real forms of hermitian symmetric spaces*, Bull. Amer. Math. Soc., 81(1975), 456-458.
- [11] H. A. Jaffee, *Anti-holomorphic automorphisms of the exceptional symmetric domains*, J. Differential Geom., 13(1978), no.1, 79-86.
- [12] S.-J. Kan, *On the characterization of Grauert tubes covered by the ball*, Math. Ann., 309 (1997), 71-80.
- [13] ———, *On the rigidity of non-positively curved Grauert tubes*, Math. Z., 229 (1998), 349-363.
- [14] ——— and D. Ma, *On the rigidity of Grauert tubes over Riemannian manifolds of constant curvature*, to appear.
- [15] ———, *On the rigidity of Grauert tubes over locally symmetric spaces*, J. Reine Angew. Math., 524(2000), 205-225.
- [16] S. Kobayashi, *Hyperbolic manifolds and holomorphic mappings*, Marcel Dekker, New York, 1970.
- [17] L. Lempert and R. Szöke, *Global solutions of the homogeneous complex Monge-Ampère equation and complex structures on the tangent bundle of Riemannian manifolds*, Math. Ann., 290 (1991), 689-712.

- [18] K.-H. Neeb, *On the complex geometry of invariant domains in complexified symmetric spaces*, Ann. Inst. Fourier, Grenoble, 49(1999), no. 1, 177-225.
- [19] G. Patrizio and P-M. Wong, *Stein manifolds with compact symmetric center*, Math. Ann., 289(1991), no.3, 355-382.
- [20] N. Sibony, *A class of hyperbolic manifolds*, in the book Recent Developments in SCV, Princeton, Princeton University Press, 1981.
- [21] K. Stein, *Überlagerungen holomorph-vollständiger komplexer Räume*, Arch. Math., 7(1956), 354-361.
- [22] R. Szöke, *Complex structures on the tangent bundles of Riemannian manifolds*, Math. Ann., 291 (1991), 409-428.
- [23] ———, *Automorphisms of certain Stein manifolds*, Math. Z., 219 (1995), 357-385.
- [24] WOLF, J. A. *Fine Structure of Hermitian Symmetric Spaces* Symmetric Spaces, ed. W. M. Boothby, G. L. Weiss; Marcel Dekker, New York 1972.
- [25] ———, *Exhaustion functions and cohomology vanishing theorems for open orbits on complex flag manifolds*, Math. Res. Letters, 2 (1995), 179-191.
- [26] ———, *Complex geometry and representations of Lie groups*, to appear in Contemporary Mathematics (Proceedings of the Bilbao Meeting in Memory of Alfred Gray).
- [27] ——— and R. Zierau, *Linear cycle spaces in flag domains*, Math. Annalen, 316 (2000), 529-545.